

CONTINUOUS SUBJECTIVE EXPECTED UTILITY WITH NON-ADDITIVE PROBABILITIES*

Peter WAKKER

*Netherlands Central Bureau of Statistics, 2270, AZ Voorburg, The Netherlands***

Submitted July 1987, accepted May 1988

A well-known theorem of Debreu about additive representations of preferences is applied in a non-additive context, to characterize continuous subjective expected utility maximization for the case where the probability measures may be non-additive. The approach of this paper does not need the assumption that lotteries with known (objective) probability distributions over consequences are available.

1. Introduction

There is nowadays an increasing interest for the approach to decision making under uncertainty which explains violations of Savage's (1954) 'sure-thing principle' by allowing for non-additivity of probability measures. A characterization of non-additivity of probability measures has been provided by Schmeidler (1984a, b, 1986). He assumed that not only consequences are available as result from a decision situation, but also all simple *lotteries* (with known probability distributions) over consequences. In many economic applications the introduction of lotteries with known probability distributions on consequences will be felt as an artificial construct. This motivated the present paper; it will remove the assumption of the availability of lotteries, and will replace that by the assumption of continuity of utility, an assumption which for most economic applications will not be felt as a restriction.

As a price for the greater generality obtained, the main characterizing condition, 'comonotonic cardinal coordinate independence', is not so simple and appealing as Schmeidler's main characterizing condition, 'comonotonic independence'. (In section 10 we shall sketch how the conditions in

*The views expressed in this paper are those of the author and do not necessarily reflect the policies of the Netherlands Central Bureau of Statistics. This paper is a rewritten version of Chapter VI of Wakker (1986). Part of the research, described in this paper, was done during a stay at the Tel Aviv University, Department of Economics, with financial support from the Netherlands Organization for the Advancement of Pure Research (Z.W.O.). The research topic of this paper was suggested to the author by D. Schmeidler. An anonymous referee gave helpful comments.

**Present address: Dept. of Mathematical Psychology, University of Nijmegen.

Schmeidler's set-up, because of the presence of lotteries, do imply comonotonic cardinal coordinate independence.) A further price for the greater generality is that, if the main theorem is just 'one-step-more-complicated' than the result of Schmeidler, the proof of the main theorem will be far more complicated. The reason is that Schmeidler could use the lotteries on the set of certain acts to obtain immediately, through certainty equivalents, a representing function unique up to a positive affine transformation. This representing function was then characterized as the Choquet integral. In our set-up there is no easy way to obtain a representing function, unique up to a positive affine transformation; obtaining it will be the major complication in our proof. A restriction of this paper is that we only consider finite state spaces. The extension to arbitrary state spaces will be given in Wakker (1987). There also many applications of Schmeidler's approach will be given.

Schmeidler extends Anscombe and Aumann's (1963) derivation of subjective expected utility to non-additive probabilities. Analogously this paper extends Wakker (1984). Gilboa (1987) analogously extends Savage (1954).

The method of proof adopted in this paper shows how a well-known theorem of Debreu (1960), characterizing additive representations of preferences, can be of use in non-additive contexts. The techniques in section 7 may also be of use for the application of Debreu's theorem to subsets of Cartesian products. Some details in the proof have not been elaborated here; they can be found in Wakker (1986).

The expression 'non-additive probability' is used in an informal way in this paper; the formal term will be 'capacity', see Definition 2.1.

Let us finally note that the main result, Theorem 5.1, can be understood without consultation of the remainder of the text, with the exception of some definitions, listed directly above that theorem.

2. The preference relation

By $S = \{1, \dots, n\}$ we denote the set of *states (of nature)*. Γ denotes a set of *consequences*. Throughout this paper we assume that Γ is a connected separable topological space. Acts are elements of the Cartesian product Γ^n , or, equivalently, functions from the set of states to the set of consequences, with act f assigning consequence f_i to state i for all i . The Cartesian product Γ^n is endowed with the product topology. By \succsim the *preference relation* of a decision maker on the set of acts is denoted. For any α in Γ , $\bar{\alpha}$ is the constant act assigning consequence α to every state. For $A \subset S$, and $f \in \Gamma^n$, f_A denotes the restriction of f to A . This can be considered to be an element of $\prod_{i \in A} \Gamma$. For $g \in \Gamma^n$, $f_{-A}g_A$ is $f_A \circ g_A$, i.e., the act which equals g on A , and f on A^c . Analogous notations are, for $1 \leq i \neq j \leq n$, $\alpha, \beta \in \Gamma$, $f \in \Gamma^n$:

$f_{-i}\alpha$ is [f with f_i replaced by α],

and

$f_{-i,j}\alpha, \beta$ is [f with f_i replaced by α and f_j by β].

The binary relation \succcurlyeq is a *weak order* on a subset E of Γ^n if it is *complete* on E (i.e., for all f, g in E : $f \succcurlyeq g$ or $g \succcurlyeq f$) and *transitive* on E (i.e., for all f, g, h in E : if $f \succcurlyeq g$ and $g \succcurlyeq h$ then $f \succcurlyeq h$). We write $f \succ g$ if $f \succcurlyeq g$ and not $g \succcurlyeq f$, $f \approx g$ if $f \succcurlyeq g$ and $g \succcurlyeq f$, $f \preccurlyeq g$ if $g \succcurlyeq f$, and $f \prec g$ if $g \succ f$. We call \succcurlyeq *trivial* on $E \subset \Gamma^n$ if $f \approx g$ for all f, g in E . By \succcurlyeq we also denote the binary relation on Γ defined by $\alpha \succcurlyeq \beta$ if $\bar{\alpha} \succcurlyeq \bar{\beta}$. In the terminologies above, and below, ‘on Γ^n ’ will often be omitted. Further \succcurlyeq is *continuous* if $\{f \in \Gamma^n: f \succcurlyeq g\}$ and $\{f \in \Gamma^n: g \succcurlyeq f\}$ are closed for all $g \in \Gamma^n$.

Let again $E \subset \Gamma^n$. A function $\phi: E \rightarrow \mathbb{R}$ *represents* \succcurlyeq on E if, for all f, g in E , $f \succcurlyeq g$ iff $\phi(f) \geq \phi(g)$. We use the term ‘increasing’ rather than ‘strictly increasing’. A *positive affine* transformation adds a real number, next multiplies with a positive real number.

Definition 2.1. A function $v: 2^S \rightarrow \mathbb{R}$ is a *capacity* if

$$v(\emptyset) = 0, \tag{2.1}$$

$$v(S) = 1, \tag{2.2}$$

$$A \subset B \Rightarrow v(A) \leq v(B) \text{ (monotonicity)}. \tag{2.3}$$

Definition 2.2. Let $v: 2^S \rightarrow \mathbb{R}$ be a capacity. Then, for any function $\phi: S \rightarrow \mathbb{R}$, the *Choquet integral* of ϕ with respect to v , denoted as $\int_S \phi dv$, or as $\int \phi dv$, is

$$\int_0^\infty v(\{i \in S: \phi(i) \geq \tau\}) d\tau + \int_{-\infty}^0 [v(\{i \in S: \phi(i) \geq \tau\}) - 1] d\tau. \tag{2.4}$$

For additive capacities the Choquet integral coincides with the usual integral, as follows from integration by parts. Let π be a permutation on $\{1, \dots, n\}$ such that

$$\phi(\pi(1)) \geq \phi(\pi(2)) \geq \dots \geq \phi(\pi(n)), \tag{2.5}$$

and write for all $1 \leq j \leq n$,

$$P^n(j) := v(\{i \in S: \pi^{-1}(i) \leq \pi^{-1}(j)\}) - v(\{i \in S: \pi^{-1}(i) < \pi^{-1}(j)\}). \tag{2.6}$$

Now it can be seen that (2.4) equals

$$\sum_{j=1}^n P^\pi(j)\phi(j). \quad (2.7)$$

We have

$$\int \lambda \phi \, dv = \lambda \int \phi \, dv \quad \text{for all } \lambda \geq 0 \quad (\text{positive homogeneity}), \quad (2.8)$$

$$\int (\lambda + \phi) \, dv = \lambda + \int \phi \, dv \quad \text{for all } \lambda \in \mathbb{R} \quad (\text{translation invariance}). \quad (2.9)$$

Further,

$$\text{if } \phi(i) \geq \phi^*(i) \text{ for all } i, \text{ then } \int \phi \, dv \geq \int \phi^* \, dv \quad (\text{monotonicity}). \quad (2.10)$$

If we consider the Choquet integral as a function(al) from \mathbb{R}^n to \mathbb{R} , with $(\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n$ interpreted as the function assigning λ_j to every j , then:

Proposition 2.1. The Choquet-integral is continuous.

Proof. See Schmeidler (1984a), or Wakker (1986, Proposition VI.2.4). \square

We need the following (2.11) in the proof of Theorem 8.1. Note that the $P^\pi(j)$ of (2.6) uniquely determine v : for any $A \subset S$, take a π such that $A = \{\pi(1), \dots, \pi(i)\}$; then $v(A) = \sum_{j=1}^i P^\pi(\pi(j))$. This also shows that if one takes an arbitrary collection of real numbers $P^\pi(j)$, one $P^\pi(j)$ for every $1 \leq j \leq n$ and permutation π on $\{1, \dots, n\}$, then there exists a (necessarily unique) capacity v such that any $P^\pi(j)$ can be derived from v as in (2.6) if and only if for all i, π, π' :

$$P^\pi(i) \geq 0; \quad \sum_{j=1}^n P^\pi(j) = 1; \quad (2.11)$$

$$[\{\pi(1), \dots, \pi(i)\} = \{\pi'(1), \dots, \pi'(i)\}] \Rightarrow \left[\sum_{j=1}^i P^\pi(\pi(j)) = \sum_{j=1}^i P^{\pi'}(\pi'(j)) \right].$$

3. Comonotonicity

The following is a central notion:

Definition 3.1. A set $C \subset \Gamma^n$ is *comonotonic* if no $f, g \in C$, $i, j \in S$ exist such that simultaneously $f_i \succ f_j$ and $g_j \succ g_i$.

Definition 3.2. For a permutation π on S , $C^\pi := \{f \in \Gamma^n: f_{\pi(1)} \succcurlyeq f_{\pi(2)} \succcurlyeq \dots \succcurlyeq f_{\pi(n)}\}$. $C^{id} = C^\pi$ with π identity.

For $f \in \Gamma^n$, \succsim_f is the binary relation on S defined by $i \succsim_f j$ whenever $f_i \succsim f_j$. If \succsim is a weak order, then so is \succsim_f . Further, for $S \subset \Gamma^n$, $\succsim_S := \bigcap_{f \in S} \succsim_f$. Thus $i \succsim_S j$ iff $f_i \succsim f_j$ for all $f \in S$. With an ordering a weak order for which no different elements are equivalent, we obtain

Lemma 3.1. Let $S \subset \Gamma^n$. Let \succsim be a weak order. The following three statements are equivalent:

- (i) S is comonotonic.
- (ii) \succsim_S is a weak order.
- (iii) $S \subset C^\pi$ for some permutation π on S .

Proof. (iii) follows from (ii) by letting π be such that $i \succ_S j \Rightarrow \pi^{-1}(i) < \pi^{-1}(j)$. If (iii) holds, then for $f, g \in S$, $f_i > f_j$ and $g_j > g_i$ would imply $\pi^{-1}(i) < \pi(j)$ and $\pi^{-1}(j) < \pi^{-1}(i)$, which cannot hold. So (i) follows. Finally, (i) is assumed, and (ii) is derived. Transitivity of \succsim_S is by transitivity of \succsim . So completeness of \succsim_S remains to be derived. If not $j \succ_S i$, then there is $f \in S$ with $f_i > f_j$. By comonotonicity $g_i \succsim_S g_j$ for all $g \in S$. So $i \succ_S j$. \square

Statements (ii) and (iii) in Lemma 3.1 were observed by Schmeidler. Because of these results he called the acts in S ‘common monotonic’, subsequently abbreviated to ‘comonotonic’. Dellacherie (1970) may have been the first to see the importance of comonotonicity in connection with the Choquet integral.

Definition 3.3. Let $C \subset \Gamma^n$. Then state i is *inessential* (with respect to \succsim) on C if $h_{-i}\alpha \approx h_{-i}\beta$ for all $h_{-i}\alpha, h_{-i}\beta \in C$. If i is inessential on C^π , then we also call i π -inessential. If π is the identity, we write *id-inessential*. The opposite of ‘inessential’ always is *essential*.

A preparatory notation:

For $\alpha, \beta \in \Gamma$, $\alpha \vee \beta$ [respectively $\alpha \wedge \beta$] is α if $\alpha \succ \beta$
[respectively $\alpha \preccurlyeq \beta$], β otherwise.

Lemma 3.2. Let \succsim be a weak order. Let $f, g \in C^\pi$, and $f_j = g_j$ for all π -essential j . Then $f \approx g$.

Proof. Suppose $f, g \in C^{\text{id}}$. Define $f^0 := f$, $g^0 := g$, and inductively, for $j = 1, \dots, n$, $f^j := f^{j-1} \vee g_j$, $g^j := g^{j-1} \vee f_j$. Note that, for all id-essential j , $f_j = g_j = f_j \vee g_j$. Note also that, for all $j > 1$, $f_j \vee g_j \preccurlyeq f_{j-1}^{j-1}$, $f_j \vee g_j \preccurlyeq g_{j-1}^{j-1}$ so that $f^j, g^j \in C^{\text{id}}$ for all j . We conclude: $f = f^0 \approx f^1 \approx \dots \approx f^n = g^n \approx g^{n-1} \approx \dots \approx g^0 = g$. \square

Lemma 3.3. Let \succcurlyeq be a weak order. Let i be π -inessential for all i, π . Then \succcurlyeq is trivial.

Proof. Let $f, g \in \Gamma^n$. Take any $\alpha \in C$. Since $\bar{\alpha} \in C^\pi$ for all π , there are π, π' such that $f, \bar{\alpha} \in C^\pi$ and $g, \bar{\alpha} \in C^{\pi'}$. By the previous Lemma, $f \approx \bar{\alpha} \approx g$. \square

4. Comonotonic cardinal coordinate independence

The following definition adapts the condition of ‘cardinal coordinate independence’ of Wakker (1984) to the present non-additive context.

Definition 4.1. The binary relation \succcurlyeq satisfies *comonotonic cardinal coordinate independence (Com.CCI)* if, for all permutations π, π' on $\{1, \dots, n\}$, all j and π -essential i , and all $f_{-i}\alpha, g_{-i}\beta, f_{-i}\gamma, g_{-i}\delta \in C^\pi$, and finally all $s_{-j}\alpha, t_{-j}\beta, s_{-j}\gamma, t_{-j}\delta \in C^{\pi'}$:

$$[f_{-i}\alpha \preceq g_{-i}\beta \ \& \ f_{-i}\gamma \succcurlyeq g_{-i}\delta \ \& \ s_{-j}\alpha \succcurlyeq t_{-j}\beta] \Rightarrow [s_{-j}\gamma \succcurlyeq t_{-j}\delta].$$

A way to gain insight into the condition is to consider the simple and elaborated proof in section 5, which may reveal the idea of strength of preference underlying the definition. Lemmas 4.1 and 4.3 will derive direct consequences of Com.CCI.

Assumption 4.1 (for this section). *The binary relation \succcurlyeq is a weak order that satisfies Com.CCI.*

Definition 4.2. The binary relation \succcurlyeq is *coordinate independent (CI)* if, for all i, f, g, α, β ,

$$[f_{-i}\alpha \succcurlyeq g_{-i}\alpha] \Leftrightarrow [f_{-i}\beta \succcurlyeq g_{-i}\beta].$$

It straightforwardly follows that, for weak orders, CI implies $[f_{-A} s_A \succcurlyeq g_{-A} s_A] \Leftrightarrow [f_{-A} t_A \succcurlyeq g_{-A} t_A]$. This we adapt to the present non-additive context, by restricting it to comonotonic acts.

Definition 4.3. The binary relation \succcurlyeq satisfies *comonotonic coordinate independence (Com.CI)* if for all comonotonic $\{f_{-A} s_A, g_{-A} s_A, f_{-A} t_A, g_{-A} t_A\}$ we have $[f_{-A} s_A \succcurlyeq g_{-A} s_A] \Leftrightarrow [f_{-A} t_A \succcurlyeq g_{-A} t_A]$.

Lemma 4.1. *The binary relation \succcurlyeq satisfies comonotonic coordinate independence.*

Proof. First we consider the special case that A contains one element, say $A = \{k\}$. Let $f_{-k} s_k, g_{-k} s_k, f_{-k} t_k, g_{-k} t_k \in C^\pi$. If k is π -inessential, then $f_{-k} s_k \approx f_{-k} t_k$, and $g_{-k} s_k \approx g_{-k} t_k$, and everything follows. So let k be π -

essential. Then $[f_{-k} s_k \preceq f_{-k} s_k \ \& \ f_{-k} t_k \succcurlyeq f_{-k} t_k \ \& \ f_{-k} s_k \succcurlyeq g_{-k} s_k]$ by Com.CCI imply $f_{-k} t_k \succcurlyeq g_{-k} t_k$.

Next we consider the general case. Say $f_{-A} s_A, g_{-A} s_A, f_{-A} t_A, g_{-A} t_A \in C^{id}$. Define:

$$a^0 := f_{-A} s_A, \quad b^0 := g_{-A} s_A, \quad c^0 := f_{-A} t_A, \quad d^0 := g_{-A} t_A.$$

Then define, inductively, for $j=1, \dots, n$:

$$\text{If } j \notin A, \text{ then } (a^j, b^j, c^j, d^j) := (a^{j-1}, b^{j-1}, c^{j-1}, d^{j-1}).$$

If $j \in A$, then set $\alpha = s_j \vee t_j$ and define

$$(a^j, b^j, c^j, d^j) := (a_{-j}^{j-1} \alpha, b_{-j}^{j-1} \alpha, c_{-j}^{j-1} \alpha, d_{-j}^{j-1} \alpha).$$

The above construction has been such that $a_k^j = c_k^j$ and $b_k^j = d_k^j$ for all $k \leq j$, and such that all new acts are in C^{id} . For instance, if $j \in A$, then $a^{j-1}, b^{j-1}, c^{j-1}, d^{j-1} \in C^{id}$ imply, by simple manipulations, $\alpha \preceq a_{j-1}^j, \alpha \preceq b_{j-1}^j, \alpha \preceq c_{j-1}^j, \alpha \preceq d_{j-1}^j$. Further $a^n = c^n, b^n = d^n$. By repeated application of the already handled case where A contains exactly one element, we conclude that each of the following preferences holds if and only if its predecessor and successor hold: $f_{-A} s_A \succcurlyeq g_{-A} s_A, a^0 \succcurlyeq b^0, a^1 \succcurlyeq b^1, \dots, a^n \succcurlyeq b^n, c^n \succcurlyeq d^n, c^{n-1} \succcurlyeq d^{n-1}, \dots, c^0 \succcurlyeq d^0, f_{-A} t_A \succcurlyeq g_{-A} t_A$. \square

Definition 4.4. We write $f_A \succcurlyeq_A^\pi g_A$ if: [there exists s_{A^c} such that $f_A s_{A^c} \succcurlyeq g_A s_{A^c}$, and $f_A s_{A^c}, g_A s_{A^c} \in C^\pi$].

Lemma 4.2. If $f_A \succcurlyeq_A^\pi g_A$, then $f_A s_{A^c} \succcurlyeq g_A s_{A^c}$ for all s_{A^c} for which $f_A s_{A^c}, g_A s_{A^c} \in C^\pi$.

Proof. Direct from Lemma 4.1. \square

Definition 4.5. The binary relation \succcurlyeq satisfies *weak monotonicity* (w.mon.) if $f \succcurlyeq g$ whenever $f_i \succcurlyeq g_i$ for all i . It satisfies *comonotonic strong monotonicity* (com.s.mon.) if $f \succ g$ for all comonotonic $\{f, g\} \subset C^\pi$ with $f_i \succcurlyeq g_i$ for all i and $f_i \succ g_i$ for a π -essential i .

Lemma 4.3. The binary relation \succcurlyeq satisfies weak, and comonotonic strong, monotonicity.

Proof. First we derive weak monotonicity. In three steps:

Step 1. Let $g = f_{-k} \alpha$, let $\{f, g\}$ be comonotonic, say $f, g \in C^{id}$, and let $f_k \succcurlyeq \alpha$. We show that $f \succcurlyeq f_{-k} \alpha$.

Suppose $f \prec f_{-k} \alpha$. Contradiction will follow.

Define, for $j=0, \dots, n$,

$$h^j \text{ has } h_1^j = \dots = h_k^j = f_k, \quad h_{j+1}^j = \dots = h_n^j = \alpha. \quad (4.1)$$

Then all h^j are in C^{id} . By Com.CI, $f < f_{-k}\alpha$ implies $h_{-k}^k f_k < h_{-k}^k \alpha$, i.e., $h^k < h^{k-1}$. Id-essentially of k follows. Thus, by Com.CCI,

$$[h_{-k}^k \alpha \leq h_{-k}^k \alpha \ \& \ h_{-k}^k \alpha \geq h_{-k}^k f_k \ \& \ h_{-j}^j \alpha \geq h_{-j}^j \alpha] \text{ implies} \\ h_{-j}^j \alpha \geq h_{-j}^j f_k, \text{ i.e., } h^{j-1} \geq h^j, \text{ for all } j \geq 1.$$

Apparently $\bar{\alpha} = h^0 \geq h^1 \geq \dots \geq h^{k-1}$, $h^{k-1} > h^k$, $h^k \geq \dots \geq h^n = \bar{f}_k$. This, finally, contradicts $f_k \geq \alpha$. Step 1 is established.

Step 2. Let again $g = f_{-k}\alpha$, and $f_k \geq \alpha$. Now we do not assume that $\{f, g\}$ is comonotonic.

Let us assume that $f \in C^{\text{id}}$. We show that $f \geq f_{-k}\alpha$.

Let l be such that $f_i \geq \alpha$, $f_j < \alpha$ for all $j > l$. Then, by repeated application of the result of Step 1, $f \geq f_{-k} f_{k+1} \geq f_{-k} f_{k+2} \geq \dots \geq f_{-k} f_l \geq f_{-k}\alpha$, since every two consecutive acts are comonotonic (e.g., $f_{-k} f_{k+2}$ and $f_{-k} f_{k+3}$ are in C^π for a π with $\pi(k+2) = k$). Step 2 is established.

Step 3. Now let $f_i \geq g_i$ for all i ; further f and g are general. We show that $f \geq g$.

By repeated application of the above result,

$$f \geq f_{-1} g_1 \geq ((f_{-1} g_1)_{-2} g_2) \geq \dots \geq (((f_{-1} g_1)_{-2} g_2) \dots_{-n} g_n) = g.$$

Weak monotonicity is established.

Next we derive com.s.mon. Suppose that $\{f, g\}$ is comonotonic, say $\{f, g\} \subset C^{\text{id}}$, and that $f_j \geq g_j$ for all j . Let further $f_k > g_k$ for an id-essential k . To derive is $f > g$. Define:

$$h \text{ has } h_j = f_j \text{ for all } j \leq k, \quad h_j = g_j \text{ for all } j > k.$$

Both $(h =) h_{-k} f_k$ and $h_{-k} g_k$ are in C^{id} . By w.mon. $f \geq h_{-k} f_k \geq h_{-k} g_k \geq g$. It is sufficient for com.s.mon. to show that $h_{-k} f_k > h_{-k} g_k$. Suppose to the contrary that $h_{-k} f_k \leq h_{-k} g_k$. We derive a contradiction.

Define h^0, \dots, h^n as in (4.1), with $\alpha = g_k$. Since k is id-essential, by Com.CCI $[h_{-k}^k g_k \leq h_{-k}^k g_k \ \& \ h_{-k}^k g_k \geq h_{-k}^k f_k \ \& \ h_{-j}^j g_k \geq h_{-j}^j g_k]$ implies $h_{-j}^j g_k \geq h_{-j}^j f_k$, i.e., $h^{j-1} \geq h^j$, for all $j \geq 1$. So $\bar{g}_k \geq \bar{f}_k$. This contradicts $f_k > g_k$. \square

Corollary 4.1. The binary relation \geq is trivial iff $\alpha \geq \beta$ for all $\alpha, \beta \in \Gamma$.

Proof. If \geq is trivial, then $\bar{\alpha} \geq \bar{\beta}$, so $\alpha \geq \beta$, for all α, β . Next assume $\alpha \geq \beta$ for all α, β . Then for any f in any C^π , and any $\alpha \in \Gamma$, $f_i \geq \alpha_i$ for all i , and $\bar{\alpha} \in C^\pi$, hence by w.mon. $f \geq \bar{\alpha}$. Analogously $f \leq \bar{\alpha}$. So $f \approx \bar{\alpha}$. Also $f \approx \bar{\alpha} \approx g$ for all f, g, α : \geq is trivial. \square

Corollary 4.2. One π has a π -essential state iff every π has a π -essential state.

Proof. If one π has a π -essential state, then \succsim is not trivial. By Corollary 4.1 $\bar{\alpha} < \bar{\beta}$ for some $\alpha, \beta \in \Gamma$. Since $\bar{\alpha}, \bar{\beta} \in C^\pi$ for every π , Lemma 3.2 implies that every π has a π -essential state. \square

5. The main theorem

In this section we give the main theorem of this paper. After the theorem the simplest implication (i) \Rightarrow (ii) in it is proved. The proof of (ii) \Rightarrow (i), and of the uniqueness results, will be carried out in the following sections, and completed in section 9. A survey of the proof is given in section 10. The elementary definitions, needed to understand the theorem, can be found in the beginning of section 2 up to Definition 2.2. Further are needed Definitions 3.1, 3.2, 3.3, and 4.1.

Theorem 5.1 (main theorem). Let $n \in \mathbb{N}$. Let Γ be a connected separable topological space. For the binary relation \succsim on Γ^n , the following two statements are equivalent:

- (i) There exist a capacity ν on $2^{\{1, \dots, n\}}$ and a continuous function $U: \Gamma \rightarrow \mathbb{R}$ such that $f \mapsto \int (U \circ f) d\nu$ represents \succsim .
- (ii) The binary relation \succsim is a continuous weak order that satisfies Com.CCI.

The following uniqueness results hold for U, ν of (i):

If some π has two or more π -essential states, then U is unique up to a positive affine transformation, and ν is uniquely determined. (5.1)

If \succsim is not trivial, and no π has two or more π -essential states, then U is unique up to a continuous increasing transformation, and ν is uniquely determined. (5.2)

If \succsim is trivial, then U is any constant function, and ν is arbitrary. (5.3)

Proof of (i) \Rightarrow (ii) above. Suppose (i) holds. Obviously \succsim is a weak order. Further, the map $f \mapsto (U(f_1), \dots, U(f_n))$ is continuous, so is, by Proposition 2.1, the map $(U(f_1), \dots, U(f_n)) \mapsto \int (U \circ f) d\nu$. Consequently the map $f \mapsto \int (U \circ f) d\nu$ is continuous. This implies continuity of \succsim .

All that remains is Com.CCI. For this, first suppose that:

$$\begin{aligned}
 & i \text{ is } \pi\text{-essential}; \quad f_{-i}\alpha \preceq g_{-i}\beta, \quad f_{-i}\gamma \succeq g_{-i}\delta; \\
 & \{f_{-i}\alpha, g_{-i}\beta, f_{-i}\gamma, g_{-i}\delta\} \subset C^\pi.
 \end{aligned}
 \tag{5.4}$$

The two preferences give, by (2.7), with the π in (2.7) identical to our present π since $f_i \succcurlyeq f_j \Rightarrow U(f_i) \geq U(f_j)$,

$$\sum_{k \neq i} P^\pi(k)U(f_k) + P^\pi(i)U(\alpha) \leq \sum_{k \neq i} P^\pi(k)U(g_k) + P^\pi(i)U(\beta)$$

and

$$\sum_{k \neq i} P^\pi(k)U(f_k) + P^\pi(i)U(\gamma) \geq \sum_{k \neq i} P^\pi(k)U(g_k) + P^\pi(i)U(\delta).$$

These two imply

$$P^\pi(i)[U(\alpha) - U(\beta)] \leq P^\pi(i)[U(\gamma) - U(\delta)]. \quad (5.5)$$

Were $P^\pi(i) = 0$, then by (2.7) and the representation of \succcurlyeq by $f \mapsto \int (U \circ f) dv$, i would be π -inessential. So

$$P^\pi(i) > 0. \quad (5.6)$$

The last two numbered results imply

$$U(\alpha) - U(\beta) \leq U(\gamma) - U(\delta). \quad (5.7)$$

Now suppose, besides (5.4), also

$$s_{-j}\alpha \succcurlyeq t_{-j}\beta; \quad \{s_{-j}\alpha, t_{-j}\beta, s_{-j}\gamma, t_{-j}\delta\} \subset C^\pi. \quad (5.8)$$

The preference implies: $\sum_{k \neq j} P^{\pi'}(k)U(s_k) + P^{\pi'}(j)U(\alpha) \geq \sum_{k \neq j} P^{\pi'}(k)U(t_k) + P^{\pi'}(j)U(\beta)$. This, and (5.7), imply

$$\sum_{k \neq j} P^{\pi'}(k)U(s_k) + P^{\pi'}(j)U(\gamma) \geq \sum_{k \neq j} P^{\pi'}(k)U(t_k) + P^{\pi'}(j)U(\delta).$$

Or: $s_{-j}\gamma \succcurlyeq t_{-j}\delta$. This is exactly what, by Com.CCI, should follow from (5.4) and (5.8). \square

6. Preparations for the proof

Lemma 6.1. Let $E \subset \Gamma^n$ be open [respectively closed] with respect to the product topology on Γ^n . Let $A \subset S$, $h \in \Gamma^n$. Then the set $\{f_A \in \prod_{i \in A} \Gamma : f_A h_{A^c} \in E\}$ is open [respectively closed] w.r.t. the product topology on $\prod_{i \in A} \Gamma$.

Proof. Say E is open. (For closed E , take complement.) Let V be the set

as defined above. Let $g_A \in V$. There exist open E_i , for all $i \in S$, such that $g_A h_{A^c}$ is in $\prod_{i \in S} E_i$, and such that the latter is a subset of E . We see that $g_A \in \prod_{i \in A} E_i \subset V$. $\prod_{i \in A} E_i$ is an open neighbourhood of g_A within V . \square

Lemma 6.2. Let \succcurlyeq be a weak order on Γ^n , continuous with respect to the product topology. Then for all $f \in \Gamma^n$, $\{\alpha \in \Gamma: \bar{\alpha} \succ f\}$ and $\{\alpha \in \Gamma: \bar{\alpha} \prec f\}$ are open subsets of Γ .

Proof. Let $\bar{\alpha} \succ f$. Then an open neighbourhood $V \subset \Gamma^n$ of $\bar{\alpha}$ exists such that $g \succ f$ for all $g \in V$. We may assume that V is of the form $A_1 \times \dots \times A_n$, with all A_j open subsets of Γ . Now $A := \bigcap_{j=1}^n A_j$ gives an open neighbourhood of α within $\{\alpha \in \Gamma: \bar{\alpha} \succ f\}$. So the latter is open.

Analogously $\{\alpha \in \Gamma: \bar{\alpha} \prec f\}$ is open. \square

Lemma 6.3. Let π have two or more π -essential states. Let the assumptions in Theorem 5.1, and also (ii) there, hold. Then also (i) and the uniqueness results there hold. If \succcurlyeq is non-trivial, then v only assigns values 0 and 1.

Proof. If there is a π with no π -essential state, then by Lemma 3.2, for all $\alpha, \beta \in \Gamma$, $\bar{\alpha} \approx \bar{\beta}$. By Corollary 4.1, \succcurlyeq is trivial. Now (5.3), and (i), follow straightforwardly. So we assume:

$$\text{Every } \pi \text{ has exactly one } \pi\text{-essential state.} \tag{6.1}$$

The binary relation on Γ , also denoted by \succcurlyeq , obviously is a weak order. By Lemma 6.2 it is continuous. By Debreu (1964) there exists a $\phi: \Gamma \rightarrow \mathbb{R}$, representing \succcurlyeq on Γ , unique up to a continuous increasing transformation. We can set $U := \phi$, as we shall see; so any continuous increasing transform of U can be used.

Next we define v . Let $A \subset S$ be arbitrary. By non-triviality we can take some α and β such that $\alpha \succ \beta$. If $\bar{\alpha}_A \bar{\beta}_{A^c} \succ \bar{\beta}$, then we define $v(A) := 1$, otherwise $v(A) := 0$. By com.s.mon. and Lemma 3.2, $v(A) = 1$ iff for any π with $\{\pi(1), \dots, \pi(k)\} = A$, A contains the π -essential state. This shows that v is independent of the particular choice of α and β above. Also it follows that $P^\pi(j) = 0$ for all π -inessential j , and $P^\pi(j) = 1$ for the π -essential j .

Now we show that with these constructions, (i) in Theorem 5.1 holds. Let f and g be two acts. Let $f \in C^\pi$, $g \in C^{\pi'}$. Let i be the π -essential state, j the π' -essential state. Then, by Lemma 3.2, $f \approx \bar{f}_i, g \approx \bar{g}_j$. There now follows: $f \succcurlyeq g \Leftrightarrow \bar{f}_i \succcurlyeq \bar{g}_j \Leftrightarrow U(f_i) \geq U(g_j) \Leftrightarrow \sum P^\pi(k) U(f_k) \geq \sum P^{\pi'}(k) U(g_k) \Leftrightarrow \int (U \circ f) dv \geq \int (U \circ g) dv$.

Finally we derive the uniqueness result (5.2). We saw above that U can be any continuous increasing transform of ϕ . Since, obviously, U has to

represent \succsim on Γ , no other kind of U can be taken: U is unique up to a continuous increasing transformation.

For uniqueness of v we consider an arbitrary π , and show that $P^\pi(i)=0$ for all π -inessential i . Then $P^\pi(j)$ must equal 1 for the π -essential j . These values $P^\pi(\cdot)$ uniquely determine v . So let, finally, $i=\pi(k)$ be π -inessential. Let $\alpha > \beta$. Let f assign α to $\pi(1), \dots, \pi(k)$, β to $\pi(k+1), \dots, \pi(n)$. Then f and $f_{-i}\beta$ are in C^π . By π -inessentiality of i , $f \approx f_{-i}\beta$. Since $U(\alpha) > U(\beta)$, by (2.7) we obtain $P^\pi(i)=0$. \square

Definition 6.1. The binary relation \succsim satisfies *restricted solvability* if, for all i, f, g, α, γ ,

$$[f_{-i}\alpha \succsim g \succsim f_{-i}\gamma] \Rightarrow [\text{there exists } \beta \text{ such that } f_{-i}\beta \approx g].$$

Lemma 6.4. A continuous weak order \succsim satisfies restricted solvability.

Proof. See Krantz et al. (1971, section 6.12.3). \square

Definition 6.2. Let $C \subset \Gamma^n$. Let $(V_j)_{j=1}^n$ be an array of functions, each from a subset of Γ to the reals. Then $(V_j)_{j=1}^n$ are *additive value functions* (for \succsim) on C if the function $V: f \mapsto \sum_{j=1}^n V_j(f_j)$ is well-defined on C , and represents \succsim on C . If a function V on a subset C of Γ^n can be written as a sum $V: f \mapsto \sum_{j=1}^n V_j(f_j)$, then V is *additive*.

Theorem 6.1 [Debreu (1960)]. Let at least three states be essential on Γ^n . The following two statements are equivalent for the binary relation \succsim on Γ^n :

- (i) There exists a continuous additive representation of the binary relation \succsim .
- (ii) The binary relation \succsim is a continuous weak order which satisfies CI.

Furthermore the function in (i) is unique up to a positive affine transformation.

Definition 6.3. $\alpha \in \Gamma$ is *maximal* [respectively *minimal*] if $\beta > \alpha$ [respectively $\beta < \alpha$] for no $\beta \in \Gamma$.

7. Additive value functions on C^{id}

In this section we derive results for C^{id} . Of course, analogous results hold for any C^π .

Assumption 7.1 (for this section). The assumptions, and statement (ii) of Theorem 5.1 hold. There are at least two id-essential states. Further, we assume that all states are id-essential. No maximal or minimal consequences exist.

The assumption of the existence of at least two id-essential states is essential for the sequel. The assumption that all states are id-essential is made only for convenience. By Lemma 3.2 id-inessential states do not affect the preference relation on C^{id} , and may just as well be suppressed from notation. They simply get assigned additive value functions V_j^{id} that are constant, say zero. Let us, as a preparation, establish ‘certainty equivalents’.

Lemma 7.1. For every $f \in \Gamma^n$ there exists α such that $f \approx \bar{\alpha}$.

Proof. For $f \in \Gamma^n$ there exist i, j , such that $f_i \succcurlyeq f_k \succcurlyeq f_j$ for all $k \in S$. Let $V := \{\alpha \in \Gamma: \bar{\alpha} \succ f\}$, $W := \{\beta \in \Gamma: f \succ \beta\}$. Then $V \cap W = \emptyset$. V and W are open by Lemma 6.2. Now $f_i \notin W$ and $f_j \notin V$ by w.mon. From connectedness of Γ existence can be derived of an $\alpha \notin V \cup W$; $\bar{\alpha} \approx f$. \square

7.1. Additive value functions $(V^h)_{j=1}^n$ on the sets E^h

Notation 7.1. For $h \in C^{\text{id}}$, $E^h := E_1^h \times \cdots \times E_n^h$ with $E_1^h := \{\alpha \in \Gamma: \alpha \succcurlyeq h_1\}$, $E_n^h := \{\alpha \in \Gamma: h_{n-1} \succcurlyeq \alpha\}$, and for all $1 \neq j \neq n$, $E_j^h := \{\alpha \in \Gamma: h_{j-1} \succcurlyeq \alpha \succcurlyeq h_j\}$.

We have $h \in E^h \subset C^{\text{id}}$. The E^h s are Cartesian products, and they are comonotonic so that on them the conditions of this paper hold without the comonotonicity premise. That enables us to apply, after some preparatory topological work, Theorem 3 of Debreu (1960).

Notation 7.2. The topology on Γ is denoted as τ . By $\tau(\succcurlyeq)$ we denote the coarsest topology on Γ with respect to which \succcurlyeq on Γ is continuous. By $\dots|E$ we denote: ‘restricted to E ’.

By Lemma 6.2 we see that $\tau(\succcurlyeq)$ is coarser than τ , so is connected and separable too. First let us note that:

Lemma 7.2. Any $E \subset \Gamma$ of the form $\{\alpha \in \Gamma: \sigma \succcurlyeq \alpha \succcurlyeq \tau\}$, $\{\alpha \in \Gamma: \sigma \succcurlyeq \alpha \succ \tau\}$, $\{\alpha \in \Gamma: \sigma \succ \alpha \succcurlyeq \tau\}$, $\{\alpha \in \Gamma: \sigma \succ \alpha \succ \tau\}$, $\{\alpha \in \Gamma: \sigma \succcurlyeq \alpha\}$, $\{\alpha \in \Gamma: \sigma \succ \alpha\}$, $\{\alpha \in \Gamma: \alpha \succcurlyeq \tau\}$, or $\{\alpha \in \Gamma: \alpha \succ \tau\}$, is connected with respect to $\tau(\succcurlyeq)|E$.

Proof. Throughout this proof, ‘open’ always refers to $\tau(\succcurlyeq)$. Let E have a form as above. Let F_1, F_2 be open in Γ . Let $E_1 = E \cap F_1$, $E_2 = E \cap F_2$. Suppose $E_1 \neq \emptyset \neq E_2$, $E_1 \cap E_2 = \emptyset$, $E_1 \cup E_2 = E$. We derive a contradiction.

Let $\alpha_1 \in E_1$, $\alpha_2 \in E_2$. $\tau(\succcurlyeq)$ does not separate between \approx -equivalent consequences, so $\alpha_1 \approx \alpha_2$ does not hold. Say $\alpha_1 \prec \alpha_2$. Define:

$$G_1 := [F_1 \cap \{\alpha: \alpha_1 \prec \alpha \prec \alpha_2\}] \cup [\{\alpha: \alpha \preccurlyeq \alpha_1\}], \text{ and}$$

$$G_2 := [F_2 \cap \{\alpha: \alpha_1 < \alpha < \alpha_2\}] \cup [\{\alpha: \alpha \succcurlyeq \alpha_2\}].$$

Then $G_1 \cap G_2 = \phi$, $G_1 \neq \phi \neq G_2$, and $G_1 \cup G_2 = \Gamma$ since $\{\alpha: \alpha_1 < \alpha < \alpha_2\} \subset E$.

First we derive openness of G_1 . For any element of G_1 , an open neighbourhood H of it within G_1 must be found. Let $\delta \in G_1$. If $\delta < \alpha_1$, take $H = \{\alpha: \alpha < \alpha_1\}$; if $\delta > \alpha_1$, $H = F_1 \cap \{\alpha: \alpha_1 < \alpha < \alpha_2\}$ is taken. So finally let $\delta \approx \alpha_1$. There must be an open neighbourhood H' of δ within F_1 of the form $\{\alpha: \alpha > \mu\}$, or $\{\alpha: v > \alpha > \mu\}$, or $\{\alpha: v > \alpha\}$ for some $\mu, v \in \Gamma$. The first case is impossible since $\alpha_2 \notin F_1$. So, finally, $H = \{\alpha: v > \alpha\}$ can be taken, in both remaining cases.

Analogously openness of G_2 is derived. Openness of G_1 and G_2 contradicts connectedness of Γ . \square

The above lemma shows that, if we use $\tau(\succcurlyeq)$ instead of τ , then every E_j^h is connected.

Lemma 7.3. Every E_j^h is separable w.r.t. $\tau(\succcurlyeq)|E_j^h$.

Proof. This is obvious if E_j^h contains exactly one \approx -equivalence class. If it contains more, then the intersection with E_j^h of any countable dense subset of Γ is dense in E_j^h , since every subset H of E_j^h , open w.r.t. $\tau(\succcurlyeq)|E_j^h$, has open, and by connectedness non-empty, intersection with $H \cap \{\alpha: h_{j-1} > \alpha > h_j\}$. \square

Lemma 7.4. For any $h \in C^{\text{id}}$, \succcurlyeq , restricted to E^h , is continuous with respect to the product topology of the $\tau(\succcurlyeq)|E_j^h$'s.

Proof. Let $f, g \in E^h$, $f > g$. We construct an auxiliary f^* such that $f^* > g$, and by means of this a subset $F_1 \times \cdots \times F_n$ of $\{v \in E^h: v > g\}$, containing f , and with every $F_j \subset E_j^h$ open with respect to $\tau(\succcurlyeq)|E_j^h$. For the construction of f_1^* , consider

$$V := \{\alpha \in \Gamma: (\alpha, f_2, \dots, f_n) > g\}.$$

By Lemma 6.1 this is open with respect to τ . V contains f_1 so is non-empty. If V contains h_1 , then $f_1^* = h_1$ and $F_1 = E_1^h$ is taken. If V does not contain h_1 , then by connectedness of Γ with respect to τ , V cannot be closed with respect to τ , so not of the form $\{\alpha: \alpha \succcurlyeq f_1\}$, as follows from Lemma 6.2. Since V , by w.mon., contains all $\alpha \succcurlyeq f_1$, V must contain an $\alpha < f_1$. This α cannot be $\leq h_1$ (that, by w.mon., would imply $h_1 \in V$). So $h_1 < \alpha < f_1: \alpha \in E_1^h$. Take $f_1^* = \alpha$, $F_1 = E_1^h \cap \{\beta \in \Gamma: \beta > \alpha\}$.

Anyway, $(f_1^*, f_2, \dots, f_n) > g$, and F_1 is open with respect to $\tau(\succcurlyeq)|E_1^h$. By

analogous constructions we obtain $f_2^*, F_2, \dots, f_n^*, F_n$, such that $(f_1^*, f_2^*, \dots, f_j^*, f_{j+1}, \dots, f_n)$ $\succ g$ for all j , $F_j = E_j^h$ if $f_j^* = h_j$, otherwise $h_j \prec f_j^* \prec f_j$ and $F_j = E_j^h \cap \{\beta: \beta \succ f_j^*\}$. Finally, $(f_1^*, \dots, f_n^*) \succ g$. For every $w \in F_1 \times \dots \times F_n$, in particular $w = f$, $w_j \succcurlyeq f_j^*$ for all j . By $w.mon.: w \succcurlyeq f^* \succ g$.

So indeed, if $f \succ g$, we can construct $F_1 \times \dots \times F_n \subset [E^h \cap \{w: w \succ g\}]$, containing f , and open with respect to the product topology of $\tau(\succcurlyeq)|_{E_j^h}$, $j=1, \dots, n$. Hence $\{f \in E^h: f \succ g\}$ is open with respect to the latter product topology, for all $g \in E^h$. Analogously $\{f \in E^h: f \prec g\}$ is open, for all g . Continuity of \succcurlyeq with respect to the product topology of the $\tau(\succcurlyeq)|_{E_j^h}$'s follows. \square

Proposition 7.1. For any $h \in C^{id}$ there exists a continuous additive representation $V^h: f \mapsto \sum_{j=1}^n V_j^h(f_j)$ for \succcurlyeq on E^h , unique up to a positive affine transformation.

Proof. We only consider the case where three or more coordinates are essential on E^h ; for the other case see Wakker (1986, Proposition VI.7.4). Since E^h , and any subset of it, is comonotonic, Com.CI holds without the comonotonicity restriction, i.e., CI holds.

Next we take care of the needed topological conditions. On every E_j^h we take as topology $\tau(\succcurlyeq)|_{E_j^h}$. By Lemma 7.2, E_j^h is connected, by Lemma 7.3 it is separable. On E^h we take the product topology. By Lemma 7.4, \succcurlyeq on E^h is continuous with respect to this topology. By Theorem 3 of Debreu (1960) (a version of Theorem 6.1 where the factor sets are allowed to be different) there is an additive representation, unique up to a positive affine transformation, and continuous w.r.t. the product topology of $\tau(\succcurlyeq)|_{E_j^h}(j=1, \dots, h)$, so certainly w.r.t. the finer 'old' topology. \square

7.2. Fitting the functions V_j^h together on C^{id}

Our next step is to show that there exist $V_j^{id}: \Gamma \rightarrow \mathbb{R}$, $j=1, \dots, n$, such that for every h and j , V_j^h can be taken to be the restriction of V_j^{id} to E_j^h . This of course could never be done if there were $A \subset S$, and $s, t \in C^{id}$, such that $(V_j^s)_{j \in A}$ and $(V_j^t)_{j \in A}$ would be additive value functions for different binary relations on the 'common domain' $\prod_{j \in A} (E_j^s \cap E_j^t)$. By comonotonic coordinate independence (Lemma 4.1) that never happens. Both $(V_j^s)_{j \in A}$ and $(V_j^t)_{j \in A}$ are additive value functions for the binary relation \succcurlyeq_A^π with $\pi = \text{identity}$, on appropriate domains.

Lemma 7.5. There exists a continuous additive function $V: f \mapsto \sum_{j=1}^n V_j(f_j)$ on C^{id} which represents \succcurlyeq on every E^h , and which is unique up to a positive affine transformation.

Proof. On every E^h we are given an additive representation $V^h: f \mapsto \sum_{j=1}^n V_j^h(f_j)$ which is unique up to a positive affine transformation. So we may add to every V_j^h an arbitrary 'location' constant $\tau_j(h)$, and multiply the V_j^h 's by one common positive 'scale constant' $\sigma(h)$, to obtain again an additive representation. The plan in the sequel is to choose, in five stages, scales and locations such that all V_j^h 's 'fit together', i.e., are the same on common domains. They can then be considered the restriction of one array $(V_j^{\text{id}})_{j=1}^n$.

There must exist $\beta^1, \beta^0 \in \Gamma$ such that $\beta^1 > \beta^0$. We shall set $V_j^{\text{id}}(\beta^0) = 0$ for all j , and $V_1^{\text{id}}(\beta^1) = 1$. (In Assumption 8.1 the scale will be changed.)

Stage 1. Choice of scale and location on E^r with $r = \beta^0$.

Let r ('reference point') = β^0 . $E_1^r = \{\alpha: \alpha \succcurlyeq \beta^0\}$, it contains β^1 . $E_n^r = \{\alpha: \alpha \preccurlyeq \beta^0\}$. For all $1 \neq j \neq n$, $E_j^r = \{\alpha: \alpha \approx \beta^0\}$. Of course we choose scale and locations such that

$$V_j^r(\beta^0) = 0 \quad \text{for all } j, \quad V_1^r(\beta^1) = 1. \quad (7.1)$$

Stage 2. Choice of scale on every E^h , and location for every V_1^h, V_n^h .

Let $h \in C^{\text{id}}$ be arbitrary. By Com.CI, (V_1^r, V_n^r) and (V_1^h, V_n^h) are additive value functions for the same $\succcurlyeq_{(1,n)}^{\text{id}}$ on $(E_1^r \cap E_1^h) \times (E_n^r \cap E_n^h)$. Note that both 1 and n are essential on $(E_1^r \cap E_1^h) \times (E_n^r \cap E_n^h)$ with respect to $\succcurlyeq_{(1,n)}^{\text{id}}$. By Lemmas 7.2 and 7.3, $E_1^r \cap E_1^h$ and $E_n^r \cap E_n^h$ are connected and separable with respect to the restrictions of $\tau(\succcurlyeq)$, and by Proposition 7.1 we get uniqueness up to a positive affine transformation. So we can choose the scale for (V_1^h, V_n^h) , (and hence for all $(V_j^h)_{j=1}^n$) and the locations for (V_1^h, V_n^h) , such that $V_1^r = V_1^h$ on $E_1^r \cap E_1^h$, and $V_n^r = V_n^h$ on $E_n^r \cap E_n^h$. Thus we shall see, even stronger,

$$V_1^s = V_1^t \quad \text{and} \quad V_n^s = V_n^t \quad \text{on common domain for all } s, t \in C^{\text{id}}. \quad (7.2)$$

This follows since, on $(E_1^s \cap E_1^t) \times (E_n^s \cap E_n^t)$, (V_1^s, V_n^s) and (V_1^t, V_n^t) are additive value functions for the same $\succcurlyeq_{(1,n)}^{\text{id}}$, hence they can differ only with respect to their locations, and a common scale. However, for $j=1, n$, V_j^s and V_j^t coincide (with V_j^r) on $(E_j^s \cap E_j^t \cap E_j^r)$; hence they coincide on common domains.

Stage 3. Intermediate observation.

In fact, for all s, t, j , V_j^s and V_j^t now have the same scale, and differ only with respect to their location, as we shall show:

$$\begin{aligned} &\text{For all } s, t \in C^{\text{id}}, \text{ and } 1 \leq j \leq n, \text{ there exist constants } \tau_j(s, t) \\ &\text{such that on } E_j^s \cap E_j^t, \quad V_j^s = \tau_j(s, t) + V_j^t. \end{aligned} \quad (7.3)$$

For $j=1$ or $j=n$, by (7.2), in fact $\tau_j(s,t)=0$. So let $1 \neq j \neq n$. Then (V_1^s, V_j^s, V_n^s) and (V_1^t, V_j^t, V_n^t) are additive value functions for the same $\succcurlyeq_{\{1,j,n\}}^{\text{id}}$ on $(E_1^s \cap E_1^t) \times (E_j^s \cap E_j^t) \times (E_n^s \cap E_n^t)$. So they can differ only by location, and common scale. However, V_1^s and V_1^t , and V_n^s and V_n^t , coincide on their common domain (which contains more than one element). The common scales must be the same.

Stage 4. Choice of location for all V_j^h 's ($j \neq 1, n$), having β^0 in their domain.

Of course for all V_j^h 's as above we choose location such that $V_j^h(\beta^0)=0$. Now not only (7.1) to (7.3) hold, but also:

$$\begin{aligned} &\text{If } V_j^s \text{ and } V_j^t \text{ have } \beta^0 \text{ in their domain then they} \\ &\text{coincide on common domain.} \end{aligned} \tag{7.4}$$

Stage 5. Choice of location for remaining V_j^h 's.

Now let $h \in C^{\text{id}}$ and j be such that $j \neq 1, j \neq n, \beta^0 \notin E_j^h = \{\alpha: h_{j-1} \succcurlyeq \alpha \succcurlyeq h_j\}$. Say $h_j \succ \beta^0 (h_{j-1} \prec \beta^0$ is analogous). Let $r(h) \in C^{\text{id}}$ be such that $(r(h))_i = h_{j-1}$ for all $i < j$, $(r(h))_i = \beta^0$ for all $i \geq j$. Then $E_j^h \subset E_j^{r(h)} = \{\alpha: h_{j-1} \succcurlyeq \alpha \succcurlyeq \beta^0\}$. By Stage 4, $V_j^{r(h)}(\beta^0)=0$. We now choose the location of V_j^h such that $V_j^h = V_j^{r(h)}$ on E_j^h . We shall show:

$$\begin{aligned} &\text{For all } s, t \in C^{\text{id}}, 1 \leq j \leq n, V_j^s \text{ and } V_j^t \text{ coincide on} \\ &\text{common domain.} \end{aligned} \tag{7.5}$$

We check this only for the case where $1 \neq j \neq n, \beta^0$ is neither in the domain of V_j^s , nor in that of V_j^t (other cases have been dealt with before, or are analogous), and $s_j \succ \beta^0$. Here E_j^s is of the form $\{\alpha: s_{j-1} \succcurlyeq \alpha \succcurlyeq s_j\}$. For E_j^t to intersect E_j^s , we must have $t_j \succ \beta^0$. Now V_j^s and $V_j^{r(s)}$ coincide on $E_j^s \cap E_j^{r(s)}$, so do V_j^t and $V_j^{r(t)}$ on $E_j^t \cap E_j^{r(t)}$; so do, by (7.4), $V_j^{r(s)}$ and $V_j^{r(t)}$ on $E_j^{r(s)} \cap E_j^{r(t)}$. The latter contains $E_j^s \cap E_j^t$. (7.5) follows.

We can now define $(V_j^{\text{id}})_{j=1}^n$. For any $\alpha \in \Gamma$, and $1 \leq j \leq n$, we take any $h \in C^{\text{id}}$ such that $\alpha \in E_j^h; h_j = \alpha$ suffices for that. Then we define $V_j^{\text{id}}(\alpha) := V_j^h(\alpha)$. By (7.5), this does not depend on the particular choice of h ; and every V_j^h is now the restriction of V_j^{id} to E_j^h .

For continuity, let $\sup(V_j^{\text{id}}(\Gamma)) > \mu > \nu > \inf(V_j^{\text{id}}(\Gamma))$. Openness of $\{f_j: \mu > V_j^{\text{id}}(f_j) > \nu\}$ follows from Proposition 7.1, with h such that $\nu > V_j^{\text{id}}(h_j)$ and, if $j > 1, V_j^{\text{id}}(h_{j-1}) > \mu$. Continuity of V_j follows.

Finally the uniqueness result. Any $(W_j^{\text{id}})_{j=1}^n$, for which real $\tau_j, j=1, \dots, n$, and positive σ exist such that $W_j^{\text{id}} = \tau_j + \sigma V_j^{\text{id}}$ for all j , satisfy the requirements of the lemma. Conversely, if $(W_j^{\text{id}})_{j=1}^n$ satisfy all the requirements of the lemma, then so do

$$Z_j^{\text{id}} = [W_j^{\text{id}} - W_j^{\text{id}}(\beta^0)] / [W_1^{\text{id}}(\beta^1) - W_1^{\text{id}}(\beta^0)].$$

From $Z_j^{\text{id}}(\beta^0) = 0$, $Z_1^{\text{id}}(\beta^1) = 1$, and from rereading the proof, the reader can see that this uniquely determines Z_j^{id} , $Z_j^{\text{id}} = V_j^{\text{id}}$ must hold for all j . \square

Note that we may not yet conclude that $(V_j^{\text{id}})_{j=1}^n$ are additive value functions on all of C^{id} .

7.3. The functions $(V_j^{\text{id}})_{j=1}^n$ are additive value functions on C^{id}

Lemma 7.6 For all (id-essential) k : $[\alpha \succcurlyeq \beta] \Leftrightarrow [V_k^{\text{id}}(\alpha) \geq V_k^{\text{id}}(\beta)]$. Hence $[\bar{\alpha} \succcurlyeq \bar{\beta}] \Leftrightarrow [\sum V_k^{\text{id}}(\alpha) \geq \sum V_k^{\text{id}}(\beta)]$.

Proof. Let α, β, k be arbitrary. Say $\alpha \succcurlyeq \beta$. Let $f_j = \alpha$ for all $j < k$, $f_j = \beta$ for all $j \geq k$. Then $(f =)f_{-k}\beta$ and $f_{-k}\alpha \in C^{\text{id}}$, and $f_{-k}\beta$ and $f_{-k}\alpha \in E^f$. By w.mon. and Lemma 7.5,

$$[\alpha \approx \beta] \Rightarrow [\alpha \succcurlyeq \beta \text{ and } \beta \succcurlyeq \alpha] \Rightarrow [f_{-k}\alpha \approx f_{-k}\beta] \Rightarrow V_k^{\text{id}}(\alpha) = V_k^{\text{id}}(\beta).$$

By com.s.mon. and Lemma 7.5,

$$\alpha \succ \beta \Rightarrow f_{-k}\alpha \succ f_{-k}\beta \Rightarrow V_k^{\text{id}}(\alpha) > V_k^{\text{id}}(\beta).$$

Analogously,

$$\alpha < \beta \Rightarrow V_k^{\text{id}}(\alpha) < V_k^{\text{id}}(\beta).$$

All of this together implies that V_k^{id} represents \succcurlyeq on Γ . \square

Lemma 7.7. Let $f \in C^{\text{id}}$, $f \approx \bar{\alpha}$. Then $\sum_{j=1}^n V_j^{\text{id}}(f_j) = \sum_{j=1}^n V_j^{\text{id}}(\alpha)$.

Proof. The case $f_j \approx \alpha$ for all j is direct. The case $f_j > \alpha$ for some j and $f_j < \alpha$ for no j , and the case $f_j < \alpha$ for some j and $f_j > \alpha$ for no j , are excluded by com.s.mon. So suppose $j < i$ exist such that $f_j > \alpha$, $f_{j+1} \approx \dots \approx f_{i-1} \approx \alpha$, $f_i < \alpha$. We define f^0 such that $f_k^0 = f_k$ for all $f_k \not\approx \alpha$, and $f_k^0 = \alpha$ for all $f_k \approx \alpha$.

Now suppose, for some $0 \leq l \leq n-2$, $f^l \in C^{\text{id}}$ has been defined such that $f^l \approx \bar{\alpha}$, and $\sum V_k^{\text{id}}(f_k^l) = \sum V_k^{\text{id}}(f_k)$, with at least l coordinates of f^l equal to α , and no coordinate of f^l equivalent but unequal to α . If in fact f^l has $l+1$ or more coordinates equal to α , define $f^{l+1} := f^l$. If not, then, say:

$$f_m^l > \alpha, \quad f_{a+1}^l = \dots = f_{b-1}^l = \alpha, \quad f_b^l < \alpha, \quad \text{with } b = a + l + 1.$$

If now $f_{-a,b}^l \alpha, \alpha \approx \bar{\alpha}$, define $f^{l+1} := f_{-a,b}^l \alpha, \alpha$.

If $f_{-a,b}^l \alpha, \alpha < \bar{\alpha}$, define, by Lemma 6.4, $\alpha < f_a^{l+1} \leq f_a^l$ such that

$$f^{l+1} := (f_{-b,a}^l \alpha, f_a^{l+1}) \approx \bar{\alpha}.$$

If $f_{-a,b}^l \alpha, \alpha > \bar{\alpha}$, define $\alpha > f_b^{l+1} \geq f_b^l$ such that

$$f^{l+1} := (f_{-a,b}^l \alpha, f_b^{l+1}) \approx \bar{\alpha}.$$

In any case, for $h = f_{-a}^l$, both f^{l+1} and f^l are in E^h , their a th coordinate being ‘between’ f_{a-1}^l and α , their b th coordinate ‘between’ α and f_b^l . Hence by Lemma 7.5, $f^l (\approx \bar{\alpha}) \approx f^{l+1}$ implies $\sum_k V_k^{\text{id}}(f_k^{l+1}) = \sum_k V_k^{\text{id}}(f_k^l)$.

Finally we end up with $f^{n-1} \approx \bar{\alpha}$, with $n-1$ coordinates equal to α . Then by com.s.mon. the remaining coordinate of f^{n-1} must also be equivalent, so equal, to α . And

$$\sum_k V_k^{\text{id}}(f_k) = \sum_k V_k^{\text{id}}(f_k^0) = \dots = \sum_k V_k^{\text{id}}(f_k^{n-1}) = \sum_k V_k^{\text{id}}(\alpha)$$

follows. \square

Theorem 7.1. *There exists a continuous additive representation $V: f \mapsto \sum_{j=1}^n V_j(f_j)$ for \succcurlyeq on C^{id} , unique up to a positive affine transformation.*

Proof. Let $f, g \in C^{\text{id}}$ be arbitrary. Let $(V_j^{\text{id}})_{j=1}^n$ be as constructed above. Let α, β be such that $f \approx \bar{\alpha}, g \approx \bar{\beta}$ (Lemma 7.1). Then $f \succcurlyeq g$ iff $\bar{\alpha} \succcurlyeq \bar{\beta}$, which by Lemma 7.6 is iff $\sum_k V_k^{\text{id}}(\alpha) \geq \sum_k V_k^{\text{id}}(\beta)$. The latter by Lemma 7.7 holds iff $\sum_k V_k^{\text{id}}(f_k) \geq \sum_k V_k^{\text{id}}(g_k)$. \square

8. Completion of the proof of Theorem 5.1 under absence of maximal and minimal consequences

Assumption 8.1 (for this section, with Theorem 8.1 excepted). *The assumptions, and statement (ii), of Theorem 5.1 hold. There exists a π with two π -essential states, say $\pi = \text{identity}$. By m we denote an id-essential state. No maximal or minimal consequences exist. Let $\beta^1 \succ \beta^0$ be two fixed consequences. For every π with two or more π -essential states, the continuous additive representation $V^\pi: f \mapsto \sum_{j=1}^n V_j^\pi(f_j)$ of \succcurlyeq on C^π (which exists according to the previous section) is chosen such that $V_j^\pi(\beta^0) = 0$ for all j , and $\sum_{j=1}^n V_j^\pi(\beta^1) = 1$.*

Note that we have changed ‘scale’, as compared to the previous section.

There we had $V_1^{\text{id}}(\beta^1) = 1$, now $\sum_{j=1}^n V_j^{\text{id}}(\beta^1) = 1$. Note also that, at present, we may not yet conclude for different π , π' , and $f \in C^\pi$, $g \in C^{\pi'}$, that $f \succcurlyeq g \Leftrightarrow \sum_{j=1}^n V_j^\pi(f_j) \geq \sum_{j=1}^n V_j^{\pi'}(g_j)$. The only consequences of Com.CCI that we used in the previous section, i.e., comonotonic coordinate independence, weak monotonicity, and comonotonic strong monotonicity, probably do not suffice for this purpose. We shall essentially use:

Lemma 8.1. *Let there be at least two π -essential, and two π' -essential, states. Let k be π' -essential. Then for all $l \in S$, $V_l^\pi = \phi_l \circ V_k^{\pi'}$ on $C^\pi \cap C^{\pi'}$ for a constant or positive affine ϕ_l .*

Proof. Say π' is the identity. We write ϕ for ϕ_l . If l is π -inessential, then V_l^π is constant, and ϕ is the same constant. So assume:

l is π -essential.

By Lemma 7.6 (which applies to all essential k) V_l^π and V_k^{id} represent the same \succcurlyeq , hence $V_l^\pi = \phi \circ V_k^{\text{id}}$ for an increasing ϕ which is continuous by Wakker (1986, Corollary VIII.10).

First note that Com.CCI implies the same property with all preferences replaced by equivalences. This we write out in terms of additive value functions, and with $\phi \circ V_k^{\text{id}}$ for V_l^π everywhere:

for all $f_{-k}\alpha, g_{-k}\beta, f_{-k}\gamma, g_{-k}\delta \in C^{\text{id}}$, and $s_{-l}\alpha, t_{-l}\beta, s_{-l}\gamma, t_{-l}\delta \in C^\pi$,

$$\{V_k^{\text{id}}(\alpha) - V_k^{\text{id}}(\beta) = {}^{(1)}\sum_{j \neq k} [V_j^{\text{id}}(g_j) - V_j^{\text{id}}(f_j)] = {}^{(2)}V_k^{\text{id}}(\gamma) - V_k^{\text{id}}(\delta)\} \quad (8.1)$$

and

$$\phi \circ V_k^{\text{id}}(\alpha) - \phi \circ V_k^{\text{id}}(\beta) = {}^{(3)}\sum_{j \neq l} [V_j^\pi(t_j) - V_j^\pi(s_j)] \quad (8.2)$$

imply

$$\sum_{j \neq l} [V_j^\pi(t_j) - V_j^\pi(s_j)] = {}^{(4)}\phi \circ V_k^{\text{id}}(\gamma) - \phi \circ V_k^{\text{id}}(\delta). \quad (8.3)$$

Now let $V_k^{\text{id}}(\mu)$ be an arbitrary element of $\text{int}(V_k^{\text{id}}(I))$. There can be seen to be an interval S around $V_k^{\text{id}}(\mu)$, so small that for all $V_k^{\text{id}}(\alpha), V_k^{\text{id}}(\beta), V_k^{\text{id}}(\gamma)$, and $V_k^{\text{id}}(\delta) \in S$, there exist f, g such that $f_{-k}\alpha, g_{-k}\beta, f_{-k}\gamma, g_{-k}\delta$ are in C^{id} , and such that $= {}^{(1)}$ is satisfied. For this we use the existence of a state $i \neq k$ which is id-essential, so that the interval $V_i^{\text{id}}(I)$ is non-degenerate. Of course, if $i < k$, then $f_i \succcurlyeq \alpha, f_i \succcurlyeq \gamma, g_i \succcurlyeq \beta, g_i \succcurlyeq \delta$ holds. If $i > k$, the reverse has to hold. Furthermore, by continuity of ϕ , S can be taken so small that $\phi(S)$ is small

enough to guarantee existence of s and t such that $s_{-i}\alpha$, $t_{-i}\beta$, $s_{-i}\gamma$, $t_{-i}\delta$ are in C^π , and such that (3) holds.

We conclude for all $V_k^{\text{id}}(\alpha)$, $V_k^{\text{id}}(\beta)$, $V_k^{\text{id}}(\gamma)$, $V_k^{\text{id}}(\delta) \in S$,

$$\begin{aligned} V_k^{\text{id}}(\alpha) - V_k^{\text{id}}(\beta) &= V_k^{\text{id}}(\gamma) - V_k^{\text{id}}(\delta) \\ \Rightarrow \phi \circ V_k^{\text{id}}(\alpha) - \phi \circ V_k^{\text{id}}(\beta) &= \phi \circ V_k^{\text{id}}(\gamma) - \phi \circ V_k^{\text{id}}(\delta). \end{aligned} \quad (8.4)$$

This is now shown by choosing f, g, s, t as above. (8.4), only for the case where $\beta = \gamma$, already suffices to show that on S , ϕ satisfies: $\phi((\alpha^* + \delta^*)/2) = [\phi(\alpha^*) + \phi(\delta^*)]/2$. By Theorem 1 of section 2.1.4 of Aczél (1966), or by (88) of section 3.7 of Hardy et al. (1934), ϕ must be affine on S . Hence it has constant derivative everywhere on $\text{int}(V_i(\Gamma))$. Consequently ϕ must be affine on $V_i(\Gamma)$. \square

For all π with two or more π -essential states, we can, by Lemma 8.1, and the fact that all $V_j^\pi(\beta^0)$ equal 0, define $\lambda_j^\pi \in \mathbb{R}_+$ such that, with m id-essential:

$$V_j^\pi = \lambda_j^\pi V_m^{\text{id}}. \quad (8.5)$$

We define, for all these π ,

$$p_j^\pi := \lambda_j^\pi / \sum_{i=1}^n \lambda_i^{\text{id}}. \quad (8.6)$$

For π with exactly one π -essential state, say l , we define

$$p_l^\pi := 1, p_i^\pi := 0 \text{ for all } i \neq l. \quad (8.7)$$

Next we define $U: \Gamma \rightarrow \mathbb{R}$.

Definition 8.1. For all $\alpha \in \Gamma$, $U: \alpha \mapsto \sum_{j=1}^n V_j^{\text{id}}(\alpha)$.

Lemma 8.2. For all π with two or more π -essential states, and all α , $V_j^\pi(\alpha) = p_j^\pi U(\alpha)$. For all π , $\sum p_j^\pi = 1$.

Proof. Let π have two π -essential states. Then

$$\begin{aligned} p_j^\pi U(\alpha) &= [\lambda_j^\pi / \sum_{i=1}^n \lambda_i^{\text{id}}] [\sum_{i=1}^n V_i^{\text{id}}(\alpha)] \\ &= [\lambda_j^\pi / \sum_{i=1}^n \lambda_i^{\text{id}}] [\sum_{i=1}^n \lambda_i^{\text{id}} V_m^{\text{id}}(\alpha)] = V_j^\pi(\alpha). \end{aligned}$$

For such π , $\sum p_j^\pi = \sum \lambda_j^\pi / \sum \lambda_i^{\text{id}} = \sum V_j^\pi(\beta^1) / \sum V_i^{\text{id}}(\beta^1) = 1/1 = 1$.

For other π , with only one π -essential state, $[\sum p_j^\pi = 1]$ is direct. \square

Lemma 8.3. Let $f \in C^\pi$, $f \approx \bar{\alpha}$. Then $\sum p_j^\pi U(f_j) = U(\alpha)$.

Proof. If there are two or more π -essential states, then by Lemma 7.7, adapted to C^π , $\sum V_j^\pi(f_j) = \sum V_j^\pi(\alpha)$. Hence then $\sum p_j^\pi U(f_j) = \sum p_j^\pi U(\alpha) = U(\alpha)$.

If π has exactly one π -essential state, say k , then by Lemma 3.2, $f \approx \bar{f}_k$. Hence by Lemma 7.6, $U(\bar{f}_k) = U(\alpha)$, i.e., $\sum p_j^\pi U(f_j) = U(\alpha)$. \square

Lemma 8.4. Let $f \in C^\pi$, $g \in C^{\pi'}$. Then $f \succcurlyeq g \Leftrightarrow \sum_{j=1}^n p_j^\pi U(f_j) \geq \sum_{j=1}^n p_j^{\pi'} U(g_j)$.

Proof. Let (Lemma 7.1) $f \approx \bar{\alpha}$, $g \approx \bar{\beta}$. Then $f \succcurlyeq g$ iff $\bar{\alpha} \succcurlyeq \bar{\beta}$, which by Lemma 7.6 holds iff $U(\alpha) \geq U(\beta)$. By Lemma 8.3 the latter holds iff $\sum p_j^\pi U(f_j) \geq \sum p_j^{\pi'} U(g_j)$. \square

Lemma 8.5. Let $A \subset S$. Let $A = \{\pi(1), \dots, \pi(k)\} = \{\pi'(1), \dots, \pi'(k)\}$. Then $\sum_{j=1}^k p_j^\pi = \sum_{j=1}^k p_j^{\pi'}$.

Proof. Let $f_j = \beta^1$ for all $j \in A$, $f_j = \beta^0$ for all $j \notin A$. Then $f \in C^\pi$ and $f \in C^{\pi'}$. Apply the above Lemma with $g = f$. \square

The following result holds in generality without Assumption 8.1 presumed.

Theorem 8.1. Let the assumptions of Theorem 5.1 hold. Let (ii) there hold. Furthermore, let no maximal or minimal consequences exist, and let there be a π with two or more π -essential states. Then (i), and (5.1), of Theorem 5.1 hold.

Proof. According to Lemma 8.5, and (2.11), with $P^\pi(j) = p_j^\pi$ for all π, j , there exists a unique capacity v in accordance with (2.6). Note that the U in Definition 8.1 is continuous. Lemma 8.4, and (2.7) now give (i) of Theorem 5.1.

To derive (5.1), say there are two id-essential states. Then the fact that $(P^{\text{id}}(j)U)_{j=1}^n$ are additive value functions for \succcurlyeq on C^{id} , and uniqueness up to a positive affine transformation of $x \mapsto \sum (V_j^{\text{id}})(x_j)$ in Theorem 7.1, give uniqueness up to a positive affine transformation of U , and together with $[\sum P^{\text{id}}(j) = 1]$ uniquely determine $(P^{\text{id}}(j))_{j=1}^n$. Analogously $(P^\pi(j))_{j=1}^n$ are uniquely determined for any π with two or more π -essential states. If π has exactly one π -essential state k , then $P^\pi(k) = 1$ must hold, and $P^\pi(j) = 0$ for all $j \neq k$. \square

9. Maximal and/or minimal consequences

In this section we derive the implication (ii) \Rightarrow (i), and the uniqueness result (5.1) in Theorem 5.1, for the case where maximal and/or minimal consequences may exist.

Assumption 9.1 (for this section). *The assumptions of Theorem 5.1 hold. Also statement (ii) there holds. There exists a permutation π with two or more π -essential states. We assume that π is identity.*

Lemma 9.1. *Let $\alpha, \gamma \in \Gamma$ be such that $\alpha \succ \gamma$. Then there exists $\beta \in \Gamma$ such that $\alpha \succ \beta \succ \gamma$.*

Proof. $E := \{\beta : \beta \succ \gamma\}$ and $F := \{\beta : \beta \prec \alpha\}$ are open and non-empty. Their union is Γ , for if $\delta \in F^c$ then $\delta \succcurlyeq \alpha$ so $\delta \succ \gamma$. Hence by connectedness of Γ , E and F must intersect. \square

Notation 9.1. $\Gamma^* := \{\alpha \in \Gamma : \alpha \text{ is neither maximal nor minimal}\}$. $G^{\pi*} := G^\pi \cap (\Gamma^*)^\pi$.

Since $\pi = \text{id}$ has a π -essential state, there exists $\alpha \succ \beta$. By Lemma 9.1, Γ^* is non-empty, and has no ('new') maximal or minimal consequences itself.

Lemma 9.2. *If i is essential on G^π (i.e. π -essential), then it is on $G^{\pi*}$.*

Proof. Say π is identity. There exist $\alpha, \beta \in \Gamma$ such that $\alpha \succ \beta$. Twice Lemma 9.1 gives γ, δ such that $\alpha \succ \gamma \succ \delta \succ \beta$. Let $f \in C^{\text{id}}$ have $f_k = \gamma$ for all $k \leq i$, $f_k = \delta$ for all $k > i$. Then $f_{-i}\gamma, f_{-i}\delta$ are in C^{id} , even in $C^{\text{id}*}$. By com.s.mon. $f_{-i}\gamma \succ f_{-i}\delta$. \square

Next we show that, on $(\Gamma^*)^\pi$, (i) in Theorem 5.1 is satisfied.

Proposition 9.1. *There exist a capacity v , and a continuous function $U^* : \Gamma^* \rightarrow \mathbb{R}$, such that $f \mapsto \int (U^* \circ f) dv$ represents \succcurlyeq on $(\Gamma^*)^\pi$.*

Proof. By Lemma 9.1, Γ^* itself has no maximal or minimal consequences. By Lemma 9.2, essentiality of states on $(\Gamma^*)^\pi$ is as on Γ^π . The proposition now follows from Theorem 8.1, if the required topological conditions can be guaranteed. This is done analogously to subsection 7.1: $\tau(\succcurlyeq) | \Gamma^*$ is taken as topology on Γ^* . Mainly by Lemma 7.2 and an analogue of Lemma 7.3, connectedness and separability are preserved. Continuity of \succcurlyeq on $(\Gamma^*)^\pi$ with respect to the product topology of the $\tau(\succcurlyeq) | \Gamma^*$'s, differs only in details from Lemma 7.4. \square

Lemma 9.3. *If α is maximal [respectively minimal], then $U^*(\Gamma^*)$ is bounded above [respectively below].*

Proof. The proof is given only for the case of a maximal α . Let $i < j$ be two id-essential states. Let, only in this proof, $(\bar{\beta}, \bar{\gamma})$ denote the act h with $h_k = \beta$ for $k \leq i$, $h_k = \gamma$ for $k > i$, for all $\beta, \gamma \in \Gamma$. By com.s.mon., for all $\gamma \in \Gamma^*$, $\bar{\alpha} \succ (\bar{\alpha}, \bar{\gamma})$.

Let $\gamma \in \Gamma^*$ be fixed, let β (by Lemma 7.1) be such that $(\bar{\alpha}, \bar{\gamma}) \approx \bar{\beta}$ (so $\beta \in \Gamma^*$). Now for all $\mu \in \Gamma^*$ with $\mu \succcurlyeq \gamma$, $(\bar{\mu}, \bar{\gamma})$ is in $\Gamma^{\text{id}*}$, and $(\bar{\mu}, \bar{\gamma}) \prec (\bar{\alpha}, \bar{\gamma}) \approx \bar{\beta}$, so

$$v(\{1, \dots, i\})U^*(\mu) + [v(S) - v(\{1, \dots, i\})]U^*(\gamma) < U^*(\beta). \quad (9.1)$$

Since i is essential on $\Gamma^{\text{id}*}$, $v(\{1, \dots, i\})$ is positive, and (9.1) induces an upper bound for $\{U^*(\mu): \mu \in \Gamma^*, \mu \succcurlyeq \gamma\}$, thus for $U^*(\Gamma^*)$. \square

Definition 9.1. If $\alpha \in \Gamma$ is maximal, then $U(\alpha) := \sup(U^*(\Gamma^*))$. If $\alpha \in \Gamma$ is minimal, then $U(\alpha) := \inf(U^*(\Gamma^*))$. If $\alpha \in \Gamma^*$, then $U(\alpha) := U^*(\alpha)$.

As we saw above, $U(\alpha) \in \mathbb{R}$ for all α .

Notation 9.2. $\Gamma^+ := \Gamma^* \cup \{\alpha \in \Gamma: \alpha \text{ is maximal}\}$.

Lemma 9.4. For all $f \in (\Gamma^+)^n$, and $\theta \in \Gamma$ with $f \approx \bar{\theta}$, $\int(U \circ f) dv = U(\theta)$.

Proof. Say $f \in \Gamma^{\text{id}}$. By com.s.mon., θ is not minimal, so $\theta \in \Gamma^+$. If no maximal α exists, Proposition 9.1 gives the desired result. So let α be maximal. Let $0 \leq k \leq n$ be such that $f_1 \approx \alpha, \dots, f_k \approx \alpha, f_{k+1} \prec \alpha, \dots, f_n \prec \alpha$. If θ is maximal, then $\theta \approx \alpha$, and by com.s.mon. $k+1, \dots, n$ must be id-inessential. Then $\int(U \circ f) dv = U(\theta)$ follows.

There remains the most complicated case, where θ is not maximal, so, neither being minimal, is in Γ^* . First we show that $\int(U \circ f) dv \leq U(\theta)$. By w.mon., for all $\mu \in \Gamma^*$ with $(\alpha \succ) \mu \succcurlyeq f_{k+1}$,

$$(\mu, \dots, \mu, f_{k+1}, \dots, f_n) \leq \theta, \quad \text{i.e.,} \quad \int(U \circ (\mu, \dots, \mu, f_{k+1}, \dots, f_n)) dv \leq U(\theta).$$

Writing, for all $1 \leq j \leq k$, $U(f_j) = U(\alpha) = \sup\{U(\mu): \mu \in \Gamma^*, \mu \succcurlyeq f_{k+1}\}$ shows that $\int(U \circ f) dv \leq U(\theta)$.

To see that $\int(U \circ f) dv \geq U(\theta)$, we consider δ such that $\theta \succ \delta$, so $f \succ \bar{\delta}$. By standard arguments continuity of \succcurlyeq , and connectedness of Γ , imply existence of μ_k such that $f_k \succ \mu_k \succcurlyeq f_{k+1}$, and $f_{-k} \mu_k \succ \bar{\delta}$. Also, μ_{k-1} exists such that $f_{k-1} \succ \mu_{k-1} \succcurlyeq \mu_k$ and $(f_{-k-1, k} \mu_{k-1}, \mu_k) \succ \bar{\delta}$. Finally we end up with $\alpha \succ \mu_1 \succcurlyeq \mu_2 \succcurlyeq \dots \succcurlyeq \mu_k$ such that $(\mu_1, \dots, \mu_k, f_{k+1}, \dots, f_n) \succ \bar{\delta}$. Hence, for all $\mu \in \Gamma$ such that $\alpha \succ \mu \succcurlyeq \mu_1 (\succcurlyeq \dots \succcurlyeq \mu_k)$, we obtain $\int(U \circ (\mu, \dots, \mu, f_{k+1}, \dots, f_n)) dv > U(\delta)$.

Substituting, for $1 \leq j \leq k$, $U(f_j) = U(\alpha) = \sup\{U(\mu): \mu \in \Gamma^*, \mu \succcurlyeq \mu_1\}$, shows that $\int(U \circ f) dv \geq U(\delta)$. This holds for all $\delta \prec \theta$. Hence $\int(U \circ f) dv \geq U(\theta)$. \square

Lemma 9.5. The map $f \mapsto \int(U \circ f) dv$ represents \succcurlyeq on $(\Gamma^+)^n$.

Proof. First for constant acts. Suppose $\gamma \succ \delta$, with γ maximal. Then, by

Lemma 9.1, $\gamma \succ \alpha \succ \delta$ for some $\alpha \in \Gamma$. So $U(\gamma) \geq U(\alpha) > U(\delta)$ follows, the latter strict inequality follows from Proposition 9.1. All other cases of $\gamma \succ \delta \Leftrightarrow U(\gamma) \geq U(\delta)$ are straightforward.

Next let $f, g \in (\Gamma^+)^n$ be arbitrary. Let $f \approx \bar{\gamma}$, $g \approx \bar{\delta}$ (by Lemma 7.1). Then $f \succ g \Leftrightarrow \bar{\gamma} \succ \bar{\delta} \Leftrightarrow U(\gamma) \geq U(\delta) \Leftrightarrow \int (U \circ f) dv \geq \int (U \circ g) dv$, the latter by Lemma 9.4. \square

Next we must turn to $(\Gamma^+ \cup \{\alpha \in \Gamma : \alpha \text{ is minimal}\})^n = \Gamma^n$, and show that also here $f \mapsto \int (U \circ f) dv$ represents \succcurlyeq . This is very analogous to the above, elaboration is left out. We conclude that the implication (ii) \Rightarrow (i) in Theorem 5.1 is now also established if maximal and/or minimal consequences exist. For the uniqueness result (5.1) in Theorem 5.1, we must show that for maximal [respectively minimal] α no other choice for $U(\alpha)$, than $\sup(U(\Gamma^*))$ [or $\inf(U(\Gamma^*))$] can be made. This can be seen for instance from the proof of Lemma 9.4. Let $i > j$ be two id-essential states. Then, with α maximal, $f_1 = \dots = f_i = \alpha$, $\alpha \succ f_{i+1} \succcurlyeq \dots \succcurlyeq f_n$, the formula $\int (U \circ f) dv = U(\theta)$ there uniquely determines $U(\alpha)$. For minimal consequences matters are analogous.

10. Survey of the proof of Theorem 5.1, and concluding remarks

The implication (i) \Rightarrow (ii) in Theorem 5.1 has been established directly below the theorem. The proof of (ii) \Rightarrow (i) for the case where no π has two or more π -essential states, and the proof of the uniqueness results (5.2) and (5.3), have been given in Lemma 6.3. There remains the case where at least one π has two or more π -essential states. The case of no maximal or minimal consequences is handled in Theorem 8.1, the existence of maximal consequences is handled in Lemma 9.5, the general case in the final lines of section 9.

Topological separability of Γ is needed only for permutations π with exactly one π -essential state. This is shown in Wakker (1986, ch. VI).

Let us now shortly and somewhat incompletely sketch how in the set-up of Schmeidler (1984a) the conditions of ‘comonotonic independence’ and w.mon. (called ‘monotonicity’ by Schmeidler) in the presence of the other ‘usual’ conditions, imply Com.CCI. Schmeidler assumes that Γ is a set of lotteries, so that a mixture operation can be defined in the usual way. Then von Neumann–Morgenstern independence requires the implication

$$f \succ g \text{ and } 0 < \lambda \leq 1 \Rightarrow \lambda f + (1 - \lambda)h \succ \lambda g + (1 - \lambda)h$$

for any f, g, h . Schmeidler weakens this to *comonotonic independence* by requiring the implication only for comonotonic $\{f, g, h\}$, and uses mainly this to characterize maximization of a Choquet-integral. Now suppose further that $f_{-i}\alpha$, $g_{-i}\beta$, $f_{-i}\gamma$, $g_{-i}\delta$, $s_{-j}\alpha$, $t_{-j}\beta$, $s_{-j}\gamma$, $t_{-j}\delta$ are as in Definition 4.1,

with the exception that now $s_{-j}\gamma < t_{-j}\delta$, in violation of Com.CCI. Let us for simplicity suppose furthermore that all preferences in Definition 4.1 are strict (the general case is derived from this through the continuity condition in Schmeidler's set-up, and mixtures with better/worse constant acts, with weights tending to zero). A violation of w.mon. will be derived. We get

$$\frac{1}{2}(f_{-i}\alpha) + \frac{1}{2}(g_{-i}\delta) < \frac{1}{2}(g_{-i}\beta) + \frac{1}{2}(g_{-i}\delta) < \frac{1}{2}(g_{-i}\beta) + \frac{1}{2}(f_{-i}\gamma),$$

the first preference by $f_{-i}\alpha < g_{-i}\beta$ and comonotonic independence, the second preference by $f_{-i}\gamma > g_{-i}\delta$ and comonotonic independence. Further

$$\frac{1}{2}(s_{-j}\alpha) + \frac{1}{2}(t_{-j}\delta) > \frac{1}{2}(t_{-j}\beta) + \frac{1}{2}(t_{-j}\delta) > \frac{1}{2}(t_{-j}\beta) + \frac{1}{2}(s_{-j}\gamma),$$

the first preference by $s_{-j}\alpha > t_{-j}\beta$ and comonotonic independence, the second by $s_{-j}\gamma < t_{-j}\delta$ and comonotonic independence. Deleting middle mixtures in the above triples, and rewriting, we get

$$[\frac{1}{2}f + \frac{1}{2}g]_{-i}[\frac{1}{2}\alpha + \frac{1}{2}\delta] < [\frac{1}{2}f + \frac{1}{2}g]_{-i}[\frac{1}{2}\beta + \frac{1}{2}\gamma],$$

and

$$[\frac{1}{2}s + \frac{1}{2}t]_{-j}[\frac{1}{2}\alpha + \frac{1}{2}\delta] > [\frac{1}{2}s + \frac{1}{2}t]_{-j}[\frac{1}{2}\beta + \frac{1}{2}\gamma].$$

If $[\frac{1}{2}\alpha + \frac{1}{2}\delta] \succcurlyeq [\frac{1}{2}\beta + \frac{1}{2}\gamma]$, then the first preference violates w.mon, if $[\frac{1}{2}\alpha + \frac{1}{2}\delta] \preccurlyeq [\frac{1}{2}\beta + \frac{1}{2}\gamma]$ then the second preference violates w.mon.

The main result of this paper, Theorem 5.1, provides, under restrictions appropriate for economic applications, a behavioural foundation for non-additive probabilities in decision making under uncertainty. It is hoped that this will help in clarifying the usefulness of non-additive probabilities.

References

- Aczél, J., 1966, Lectures on functional equations and their applications (Academic Press, New York).
- Anscombe, F.J. and R.J. Aumann, 1963, A definition of subjective probability, *Annals of Mathematical Statistics* 34, 199–205.
- Choquet, G., 1953–54, Theory of capacities, *Annales de l'Institut Fourier (Grenoble)*, 131–295.
- Debreu, G., 1960, Topological methods in cardinal utility theory, in: K.J. Arrow, S. Karlin and P. Suppes, eds., 1959, *Mathematical methods in the social sciences* (Stanford University Press, Stanford, CA) 16–26.
- Debreu, G., 1964, Continuity properties of Paretian utility, *International Economic Review* 5, 285–293.
- Dellacherie, C., 1970, Quelques commentaires sur les prolongements de capacités, *Seminaire de Probabilités V Strasbourg, Lecture Notes in Mathematics*, no. 191 (Springer, Berlin).
- Gilboa, I., 1987, Expected utility with purely subjective non-additive probabilities, *Journal of Mathematical Economics* 16, 65–88.
- Hardy, G.H., J.E. Littlewood and G. Pólya, 1934, *Inequalities* (Cambridge University Press, Cambridge).

- Krantz, D.H., R.D. Luce, P. Suppes and A. Tversky, 1971, *Foundations of measurement, Vol. I: Additive and polynomial representations* (Academic Press, New York).
- Savage, L.J., 1954, *The foundations of statistics* (Wiley, New York).
- Schmeidler, D., 1984a, *Subjective probability and expected utility without additivity*, Caress working paper 84-21 (first part) (Center for Analytic Research in Economics and the Social Sciences, University of Pennsylvania, Philadelphia, PA).
- Schmeidler, D., 1984b, *Nonadditive probabilities and convex games*, Caress working paper 84-21 (second part) (Center for Analytic Research in Economics and the Social Sciences, University of Pennsylvania, Philadelphia, PA).
- Schmeidler, D., 1986, *Integral representation without additivity*, *Proceedings of the American Mathematical Society* 97, 255-261.
- Wakker, P.P., 1984, *Cardinal coordinate independence for expected utility*, *Journal of Mathematical Psychology* 28, 110-117.
- Wakker, P.P., 1986, *Representations of choice situations*, Ph.D. dissertation (Department of Economics, University of Brabant, Tilburg).
- Wakker, P.P., 1987, *Nonadditive probabilities and derived strengths of preferences*, Internal report 87 MA 03 (Department of Mathematical Psychology, University of Nijmegen, Nijmegen).